

Quasicompact and Riesz unital endomorphisms of real Lipschitz algebras of complex-valued functions

Maliheh Mayghani¹ and Davood Alimohammadi^{2*}

ABSTRACT. We first show that a bounded linear operator T on a real Banach space E is quasicompact (Riesz, respectively) if and only if $T' : E_{\mathbb{C}} \rightarrow E_{\mathbb{C}}$ is quasicompact (Riesz, respectively), where the complex Banach space $E_{\mathbb{C}}$ is a suitable complexification of E and T' is the complex linear operator on $E_{\mathbb{C}}$ associated with T . Next, we prove that every unital endomorphism of real Lipschitz algebras of complex-valued functions on compact metric spaces with Lipschitz involutions is a composition operator. Finally, we study some properties of quasicompact and Riesz unital endomorphisms of these algebras.

1. INTRODUCTION AND PRELIMINARIES

The symbol \mathbb{K} denotes a field that can be either \mathbb{R} or \mathbb{C} . Let E be an infinite dimensional Banach space over \mathbb{K} . We denote by $\mathcal{B}_{\mathbb{K}}(E)$ and $\mathcal{K}_{\mathbb{K}}(E)$ the set of all bounded linear operators and compact linear operators over \mathbb{K} on E , respectively. It is known that $\mathcal{B}_{\mathbb{K}}(E)$ with the operator norm is a unital Banach algebra and $\mathcal{K}_{\mathbb{K}}(E)$ is a closed ideal of $\mathcal{B}_{\mathbb{K}}(E)$ over \mathbb{K} . The essential norm $\|T\|_e$ of $T \in \mathcal{B}_{\mathbb{K}}(E)$ is the norm of $T + \mathcal{K}_{\mathbb{K}}(E)$ in the Calkin algebra $\mathcal{B}_{\mathbb{K}}(E)/\mathcal{K}_{\mathbb{K}}(E)$, i.e.,

$$\begin{aligned}\|T\|_e &= \|T + \mathcal{K}_{\mathbb{K}}(E)\| \\ &= \inf\{\|T - S\| : S \in \mathcal{K}_{\mathbb{K}}(E)\}.\end{aligned}$$

2010 *Mathematics Subject Classification.* 47B48, 46J10, 47B38.

Key words and phrases. Complexification, Lipschitz algebra, Lipschitz involution, Quasicompact operator, Riesz operator, Unital endomorphism.

Received: 22 November 2016, Accepted: 20 December 2016.

* Corresponding author.

The essential spectral radius $r_e(T)$ of $T \in \mathcal{B}_{\mathbb{K}}(E)$ is the spectral radius of $T + \mathcal{K}_{\mathbb{K}}(E)$ in $\mathcal{B}_{\mathbb{K}}(E)/\mathcal{K}_{\mathbb{K}}(E)$, i.e.,

$$\begin{aligned} r_e(T) &= \lim_{n \rightarrow \infty} (\|T^n\|_e)^{\frac{1}{n}} \\ &= \inf \left\{ (\|T^n\|_e)^{\frac{1}{n}} : n \in \mathbb{N} \right\}. \end{aligned}$$

The linear operator $T \in \mathcal{B}_{\mathbb{K}}(E)$ is called power compact if $T^N \in \mathcal{K}_{\mathbb{K}}(E)$ for some $N \in \mathbb{N}$, Riesz if $r_e(T) = 0$ and quasicompact if $r_e(T) < 1$. Clearly, $T \in \mathcal{K}_{\mathbb{K}}(E)$ if and only if $\|T\|_e = 0$, and T is quasicompact if and only if $\|T^n\|_e < 1$ for some $n \in \mathbb{N}$. Moreover, T is quasicompact if T is a Riesz operator and T is Riesz operator if T is power compact.

Let A be a unital commutative Banach algebra with unit e_A over \mathbb{K} . A linear map $T : A \rightarrow A$ is called an endomorphism of A if $T(fg) = (Tf)(Tg)$ for all $f, g \in A$. An endomorphism T of A is unital if $T(e_A) = e_A$.

Feinstein and Kamowitz studied quasicompact and Riesz unital endomorphisms of commutative semisimple unital complex Banach algebras in [5]. They investigated their studies in [6] whenever Banach algebras considered semiprime. We recall that a complex algebra A is semiprime if $J = \{0\}$ is the only ideal of A with $J^2 = \{0\}$.

Let E be a linear space over \mathbb{K} and $T : E \rightarrow E$ be a linear operator of E . We denote by $\sigma_p(T)$ and $\sigma(T)$, the set of all eigenvalues of T and the spectrum of T in \mathbb{K} , respectively, i.e.,

$$\begin{aligned} \sigma_p(T) &= \{\lambda \in \mathbb{K} : \lambda I - T \text{ is not injective}\}, \\ \sigma(T) &= \{\lambda \in \mathbb{K} : \lambda I - T \text{ is not invertible}\}. \end{aligned}$$

Let X be a compact Hausdorff space. We denote by $C_{\mathbb{K}}(X)$ the set of all \mathbb{K} -valued continuous functions on X . It is known that $C_{\mathbb{K}}(X)$ is a unital commutative Banach algebra over \mathbb{K} with unit 1_X , the constant function on X with value 1, where equipped with the uniform norm

$$\|f\|_X = \sup\{|f(x)| : x \in X\}, \quad (f \in C_{\mathbb{K}}(X)).$$

We write $C(X)$ instead of $C_{\mathbb{C}}(X)$. Let A be a unital subalgebra of $C_{\mathbb{K}}(X)$ over \mathbb{K} which is a Banach algebra under an algebra norm $\|\cdot\|$. If $\varphi : X \rightarrow X$ is a self-map of X such that $f \circ \varphi \in A$ for each $f \in A$, then the map $T : A \rightarrow A$ defined by $Tf = f \circ \varphi$ ($f \in A$), is a unital endomorphism of A which is called the composition endomorphism of A induced by φ . Recall that if $T : A \rightarrow A$ is the composition endomorphism of A induced by a self-map $\varphi : X \rightarrow X$, then $T^n : A \rightarrow A$ is the composition endomorphism of A induced by the self-map $\varphi_n : X \rightarrow X$, where φ_n is the n th iterate of φ . We also consider φ_0 the identity self-map of X .

Let (X, d) and (Y, ρ) be metric spaces such that X and Y have infinitely many points. A map $\varphi : X \rightarrow Y$ is called a Lipschitz mapping from

(X, d) into (Y, ρ) if there exists a constant C such that $\rho(\varphi(x), \varphi(y)) \leq Cd(x, y)$ for all $x, y \in X$. The constant Lipschitz of a map $\varphi : X \rightarrow Y$ is denoted by $p(\varphi)$ and defined by

$$p(\varphi) = \sup \left\{ \frac{\rho(\varphi(x), \varphi(y))}{d(x, y)} : x, y \in X, x \neq y \right\}.$$

Thus $\varphi : X \rightarrow Y$ is a Lipschitz mapping from (X, d) into (Y, ρ) if and only if $p(\varphi) < \infty$. A map $\varphi : X \rightarrow Y$ is called a supercontractive mapping from (X, d) into (Y, ρ) if

$$\lim_{d(x,y) \rightarrow 0} \frac{\rho(\varphi(x), \varphi(y))}{d(x, y)} = 0.$$

Let (X, d) be a metric space such that X has infinitely many points. A self-map $\varphi : X \rightarrow X$ is called a Lipschitz mapping on (X, d) if φ is a Lipschitz mapping from (X, d) into (X, d) . A function $f : X \rightarrow \mathbb{K}$ is called a \mathbb{K} -valued Lipschitz function on (X, d) if f is a Lipschitz mapping from (X, d) into the Euclidean metric space \mathbb{K} . We denote by $p_{(X,d)}(f)$ the constant Lipschitz of $f : X \rightarrow \mathbb{K}$, i.e.,

$$p_{(X,d)}(f) = \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)} : x, y \in X, x \neq y \right\}.$$

A function $f : X \rightarrow \mathbb{K}$ is called a \mathbb{K} -valued supercontractive function on (X, d) if f is a supercontractive mapping from (X, d) into the Euclidean space \mathbb{K} . For $\alpha \in (0, 1]$, the formula $d^\alpha(x, y) = (d(x, y))^\alpha$ defines a new metric d^α on X such that the generated topology on X by d^α coincides with the generated topology on X by d .

Let (X, d) be a compact metric space. We denote by $\text{Lip}_{\mathbb{K}}(X, d^\alpha)$ the set of all \mathbb{K} -valued Lipschitz functions on (X, d^α) . Then $\text{Lip}_{\mathbb{K}}(X, d^\alpha)$ is a subalgebra of $C_{\mathbb{K}}(X)$ over \mathbb{K} containing 1_X and separates the points of X . Moreover, $\text{Lip}_{\mathbb{K}}(X, d^\alpha)$ is a commutative unital Banach algebra over \mathbb{K} under the d^α -Lipschitz norm

$$\|f\|_{\text{Lip}_{\mathbb{K}}(X, d^\alpha)} = \|f\|_X + p_{(X, d^\alpha)}(f), \quad (f \in \text{Lip}_{\mathbb{K}}(X, d^\alpha)).$$

Note that the infiniteness of X implies that $\text{Lip}_{\mathbb{K}}(X, d^\alpha)$ is infinite dimensional. For $\alpha \in (0, 1)$, the set of all supercontractive \mathbb{K} -valued functions on (X, d^α) is denoted by $\text{lip}_{\mathbb{K}}(X, d^\alpha)$. Clearly, $\text{lip}_{\mathbb{K}}(X, d^\alpha)$ is a subalgebra of $\text{Lip}_{\mathbb{K}}(X, d^\alpha)$ over \mathbb{K} containing 1_X and closed in $(\text{Lip}_{\mathbb{K}}(X, d^\alpha), \|\cdot\|_{\text{Lip}_{\mathbb{K}}(X, d^\alpha)})$. So $(\text{lip}_{\mathbb{K}}(X, d^\alpha), \|\cdot\|_{\text{Lip}_{\mathbb{K}}(X, d^\alpha)})$ is a unital commutative Banach algebra over \mathbb{K} . It is clear that $\text{Lip}_{\mathbb{K}}(X, d^\beta) \subseteq \text{lip}_{\mathbb{K}}(X, d^\alpha)$ where $0 < \alpha < \beta \leq 1$. We write $\text{Lip}(X, d^\alpha)$ and $\text{lip}(X, d^\alpha)$ instead of $\text{Lip}_{\mathbb{C}}(X, d^\alpha)$ and $\text{lip}_{\mathbb{C}}(X, d^\alpha)$, respectively.

The algebras $\text{Lip}_{\mathbb{K}}(X, d^\alpha)$ for $\alpha \in (0, 1]$ and $\text{lip}_{\mathbb{K}}(X, d^\alpha)$ for $\alpha \in (0, 1)$ are called Lipschitz algebra and little Lipschitz algebra of order α on

(X, d) , respectively. These algebras were first introduced by Sherbert in [15, 16]. He showed [15, Theorem 5.1] that a linear map $T : \text{Lip}(X, d^\alpha) \rightarrow \text{Lip}(X, d^\alpha)$ is a unital endomorphism if and only if there exists a Lipschitz mapping $\varphi : X \rightarrow X$ on (X, d) such that T is induced by φ . One can show that this result holds for $\text{lip}(X, d^\alpha)$ instead of $\text{Lip}(X, d^\alpha)$.

Behrouzi studied quasicompact and Riesz unital endomorphisms of $\text{Lip}(X, d^\alpha)$ for $\alpha \in (0, 1]$ and $\text{lip}(X, d^\alpha)$ for $\alpha \in (0, 1)$ in [3].

Jiménez-Vargas, Lacrus and Villegas-Vallecillos in [8] studied the essential norm of composition operators on Banach spaces of Hölder functions on pointed compact metric spaces over \mathbb{K} and got a formula for it.

Golbaharan and Mahyar in [7] generalized some results of [3], studied essential spectral radius of a unital endomorphism of these algebras and got a formula for it.

Sanatpour investigated quasicompact composition operators on certain classes of complex Lipschitz algebras and obtained certain properties of power-contractive self-maps of compact plane sets in [14].

Quasicompact and Riesz endomorphisms of certain subalgebras of Lipschitz algebras were studied in [11, 12, 13].

Let X be a topological space. A self-map $\tau : X \rightarrow X$ is called a topological involution on X if τ is continuous and $\tau(\tau(x)) = x$ for all $x \in X$. Clearly, such τ is a homeomorphism from X onto X .

Let X be a compact Hausdorff space and τ be a topological involution on X . The map $\tau^* : C(X) \rightarrow C(X)$ defined by $\tau^*(f) = \bar{f} \circ \tau$ is an algebra involution on $C(X)$, which is called the algebra involution induced by τ on $C(X)$. We now define

$$C(X, \tau) = \{f \in C(X) : \tau^*(f) = f\}.$$

Then $C(X, \tau)$ is a unital self-adjoint uniformly closed real subalgebra of $C(X)$ that separates the points of X , $i1_X \notin C(X, \tau)$, $C(X) = C(X, \tau) \oplus iC(X, \tau)$ and

$$\max\{\|f\|_X, \|g\|_X\} \leq \|f + ig\|_X \leq 2 \max\{\|f\|_X, \|g\|_X\},$$

for all $f, g \in C(X, \tau)$. Real Banach algebra $C(X, \tau)$ was defined explicitly by Kulkarni and Limaye in [9]. For further general facts about $C(X, \tau)$ and certain real subalgebras, we refer to [10].

Let (X, d) be a metric space. A self-map $\tau : X \rightarrow X$ is called a Lipschitz involution on (X, d) if $\tau(\tau(x)) = x$ for all $x \in X$ and τ is a Lipschitz mapping on (X, d) .

Note that if τ is a Lipschitz involution on (X, d) , then τ is a topological involution on (X, d) and $p(\tau) \geq 1$.

Let (X, d) be a compact metric space and τ be a topological involution on (X, d) . Then for $\alpha \in (0, 1]$ we have $\tau^*(\text{Lip}(X, d^\alpha)) =$

$\text{Lip}(X, d^\alpha)$, $p_{(X, d^\alpha)}(\tau^*(f)) \leq (p(\tau))^\alpha p_{(X, d^\alpha)}(f)$ and $\|\tau^*(f)\|_{\text{Lip}(X, d^\alpha)} \leq (p(\tau))^\alpha \|f\|_{\text{Lip}(X, d^\alpha)}$ for all $f \in \text{Lip}(X, d^\alpha)$. Moreover, $\tau^*(\text{lip}(X, d^\alpha)) = \text{lip}(X, d^\alpha)$ for $\alpha \in (0, 1)$. We now define

$$\begin{aligned} \text{Lip}(X, d^\alpha, \tau) &:= \{f \in \text{Lip}(X, d^\alpha) : \tau^*(f) = f\}, \quad (0 < \alpha \leq 1), \\ \text{lip}(X, d^\alpha, \tau) &:= \{f \in \text{lip}(X, d^\alpha) : \tau^*(f) = f\}, \quad (0 < \alpha < 1). \end{aligned}$$

In fact, $\text{Lip}(X, d^\alpha, \tau) = \text{Lip}(X, d^\alpha) \cap C(X, \tau)$ and $\text{lip}(X, d^\alpha, \tau) = \text{lip}(X, d^\alpha) \cap C(X, \tau)$.

The following result is a modification of [1, Theorem 2.7].

Theorem 1.1. *Let (X, d) be a compact metric space and τ be a Lipschitz involution on (X, d) . Suppose that $A = \text{Lip}(X, d^\alpha, \tau)$ and $B = \text{Lip}(X, d^\alpha)$ for $\alpha \in (0, 1]$ ($A = \text{lip}(X, d^\alpha, \tau)$ and $B = \text{lip}(X, d^\alpha)$ for $\alpha \in (0, 1)$, respectively). Then the following statements hold.*

- (i) A is a self-adjoint real subalgebra of B , $1_X \in A$ and $i1_X \notin A$.
- (ii) $B = A \oplus iA$ and A separates the points of X .
- (iii) For all $f, g \in A$;

$$\begin{aligned} \max\{\|f\|_{\text{Lip}(X, d^\alpha)}, \|g\|_{\text{Lip}(X, d^\alpha)}\} &\leq (p(\tau))^\alpha \|f + ig\|_{\text{Lip}(X, d^\alpha)} \\ &\leq 2(p(\tau))^\alpha \max\{\|f\|_{\text{Lip}(X, d^\alpha)}, \|g\|_{\text{Lip}(X, d^\alpha)}\}. \end{aligned}$$

- (iv) A is closed in $(B, \|\cdot\|_{\text{Lip}(X, d^\alpha)})$ and so $(A, \|\cdot\|_{\text{Lip}(X, d^\alpha)})$ is a real Banach algebra.
- (v) $A = \text{Lip}_{\mathbb{R}}(X, d^\alpha)$ ($A = \text{lip}_{\mathbb{R}}(X, d^\alpha)$, respectively), if and only if τ is the identity map on X .

The algebras $\text{Lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1]$ and $\text{lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1)$ are called real Lipschitz algebra and real little Lipschitz algebra of complex-valued functions of order α on $((X, d), \tau)$, respectively. These algebras were first studied in [1]. In this paper we study some properties of quasicompact and Riesz unital endomorphisms of $\text{Lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1]$ and $\text{lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1)$, considering obtained results in [7]. Moreover, we show that the class of quasicompact (Riesz, respectively) unital endomorphisms of real Lipschitz algebras of complex-valued functions is larger than the class of quasicompact (Riesz, respectively) unital endomorphisms of complex Lipschitz algebras.

2. RESULTS

Let E be a real linear space (real algebra, respectively). A complex linear space (complex algebra, respectively) $E_{\mathbb{C}}$ is called a complexification of E if there exists an injective real linear map (real algebra homomorphism, respectively) $J : E \rightarrow E_{\mathbb{C}}$ such that $E_{\mathbb{C}} = J(E) \oplus iJ(E)$. For

example if A is a real algebra, then $A \times A$ with algebra operations

$$\begin{aligned} (a_1, b_1) + (a_2, b_2) &= (a_1 + a_2, b_1 + b_2), \quad (a_1, b_1, a_2, b_2 \in A), \\ (\alpha + i\beta)(a, b) &= (\alpha a - \beta b, \beta a + \alpha b), \quad (\alpha, \beta \in \mathbb{R}, a, b \in A), \\ (a_1, b_1)(a_2, b_2) &= (a_1 a_2 - b_1 b_2, a_1 b_2 + b_1 a_2), \quad (a_1, b_1, a_2, b_2 \in A), \end{aligned}$$

is a complexification of A when $J : A \rightarrow A \times A$ is defined by $J(a) = (a, 0)$ ($a \in A$).

Applying the proof of [4, Proposition I.1.13], one can show that if $(E, \|\cdot\|)$ is a real Banach space, then there exists a norm $\|\cdot\|$ on $E \times E$ such that $\|(a, 0)\| = \|a\|$ for each $a \in E$ and $\max\{\|a\|, \|b\|\} \leq \|(a, b)\| \leq 2 \max\{\|a\|, \|b\|\}$ for all $a, b \in E$.

Definition 2.1. Let E be a real linear space and $E_{\mathbb{C}}$ be a complexification of E under an injective real linear map $J : E \rightarrow E_{\mathbb{C}}$. Suppose that $T : E \rightarrow E$ is a real linear operator on E and the map $T' : E_{\mathbb{C}} \rightarrow E_{\mathbb{C}}$ is defined by

$$T'(J(a) + iJ(b)) = J(Ta) + iJ(Tb), \quad (a, b \in E).$$

Clearly, T' is a complex linear operator on $E_{\mathbb{C}}$. We say that T' is the complex linear operator on $E_{\mathbb{C}}$ associated with T .

The following result is a modification of [2, Theorem 2.1].

Theorem 2.2. Let $(E, \|\cdot\|)$ be a real Banach space, $E_{\mathbb{C}}$ be a complexification of E under an injective real linear map $J : E \rightarrow E_{\mathbb{C}}$. Suppose that $\|\cdot\|$ is a norm on $E_{\mathbb{C}}$ with $\|J(a)\| = \|a\|$ for each $a \in E$ and there exist positive constants k_1 and k_2 such that

$$\max\{\|a\|, \|b\|\} \leq k_1 \|J(a) + iJ(b)\| \leq k_2 \max\{\|a\|, \|b\|\},$$

for all $a, b \in E$. Let $T \in \mathcal{B}_{\mathbb{R}}(E)$ and $T' : E_{\mathbb{C}} \rightarrow E_{\mathbb{C}}$ be the complex linear operator on $E_{\mathbb{C}}$ associated with T . Then the following statements hold.

- (i) $T' \in \mathcal{B}_{\mathbb{C}}(E_{\mathbb{C}})$ and $\|T'\| \leq k_1 k_2 \|T\|$.
- (ii) T is compact if and only if T' is compact.
- (iii) T is injective if and only if T' is injective.
- (iv) T is invertible if and only if T' is invertible.
- (v) $\sigma_p(T) = \mathbb{R} \cap \sigma_p(T')$.
- (vi) $\sigma(T) = \mathbb{R} \cap \sigma(T')$.

Theorem 2.3. Let $(E, \|\cdot\|)$ be a real Banach space and $E_{\mathbb{C}}$ be a complexification of E under an injective real linear map $J : E \rightarrow E_{\mathbb{C}}$. Suppose that $\|\cdot\|$ is a norm on $E_{\mathbb{C}}$ with $\|J(a)\| = \|a\|$ for each $a \in E$ and there exist positive constants k_1 and k_2 such that

$$\max\{\|a\|, \|b\|\} \leq k_1 \|J(a) + iJ(b)\| \leq k_2 \max\{\|a\|, \|b\|\},$$

for all $a, b \in E$. Let $T \in \mathcal{B}_{\mathbb{R}}(E)$ and $T' : E_{\mathbb{C}} \rightarrow E_{\mathbb{C}}$ be the complex linear operator on $E_{\mathbb{C}}$ associated with T . Then the following statements hold.

- (i) $r_e(T') = r_e(T)$.
- (ii) T is quasicompact if and only if T' is quasicompact.
- (iii) T is Riesz if and only if T' is Riesz.

Proof. (i). Let $S \in \mathcal{K}_{\mathbb{R}}(E)$. Define the map $S' : E_{\mathbb{C}} \rightarrow E_{\mathbb{C}}$ by

$$S'(J(a) + iJ(b)) = J(Sa) + iJ(Sb) \quad (a, b \in E).$$

Then $S' \in \mathcal{K}_{\mathbb{C}}(E_{\mathbb{C}})$ and $\|S'\| \leq k_1 k_2 \|S\|$ by parts (ii) and (i) of Theorem 2.2, respectively. Moreover,

$$\|(T')^n - S'\| = \|(T^n - S)'\| \leq k_1 k_2 \|T^n - S\|,$$

for all $n \in \mathbb{N}$, by Theorem 2.2. This implies that

$$\|(T')^n\|_e \leq k_1 k_2 \|T^n\|_e,$$

for all $n \in \mathbb{N}$. Therefore,

$$(2.1) \quad r_e(T') \leq r_e(T).$$

Define the map $\Psi_1 : E \rightarrow E_{\mathbb{C}}$ by $\Psi_1(a) = J(a) + i0$, ($a \in E$) and the map $P_1 : E_{\mathbb{C}} \rightarrow E$ by $P_1(J(a) + iJ(b)) = a$, ($a, b \in E$). Clearly, Ψ_1 is a bounded real linear operator from E into $E_{\mathbb{C}}$, $\|\Psi_1\| \leq 1$, P_1 is a bounded real linear operator from $E_{\mathbb{C}}$ into E and $\|P_1\| \leq k_1$. Moreover, $P_1 \circ (T')^n \circ \Psi_1 = T^n$ for all $n \in \mathbb{N}$. Let $S \in \mathcal{K}_{\mathbb{C}}(E_{\mathbb{C}})$. Define the map $S_1 : E \rightarrow E$ by $S_1 = P_1 \circ S \circ \Psi_1$. Then $S_1 \in \mathcal{K}_{\mathbb{R}}(E)$ and

$$\begin{aligned} \|T^n - S_1\| &= \|P_1 \circ ((T')^n - S) \circ \Psi_1\| \\ &\leq \|P_1\| \|((T')^n - S)\| \|\Psi_1\| \\ &\leq k_1 \|((T')^n - S)\|, \end{aligned}$$

for all $n \in \mathbb{N}$. Hence,

$$\|T^n\|_e \leq k_1 \|(T')^n\|_e,$$

for all $n \in \mathbb{N}$. Therefore,

$$(2.2) \quad r_e(T) \leq r_e(T').$$

From (2.1) and (2.2), we get $r_e(T') = r_e(T)$ and so (i) holds. Clearly, (ii) and (iii) are hold by (i). \square

Theorem 2.4. *Let (X, d) be a compact metric space, $\tau : X \rightarrow X$ be a Lipschitz involution on (X, d) and $A = \text{Lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1]$ or $A = \text{lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1)$.*

- (i) *If $\varphi : X \rightarrow X$ is a Lipschitz mapping on (X, d) satisfying $\varphi \circ \tau = \tau \circ \varphi$, then $f \circ \varphi \in A$ for all $f \in A$.*

- (ii) *If $T : A \rightarrow A$ is a unital endomorphism of A , then T is bounded and there exists a Lipschitz mapping φ on (X, d) satisfying $\varphi \circ \tau = \tau \circ \varphi$ such that T is induced by φ .*

Proof. Let $B = \text{Lip}(X, d^\alpha)$ if $A = \text{Lip}(X, d^\alpha, \tau)$ and $B = \text{lip}(X, d^\alpha)$ if $A = \text{lip}(X, d^\alpha, \tau)$. Clearly, B is the complexification of A under the injective real algebra homomorphism $J : A \rightarrow B$ defined by $J(f) = f$ ($f \in A$). Moreover, by Theorem 1.1, we have

$$\begin{aligned} \max\{\|f\|_{\text{Lip}(X, d^\alpha)}, \|g\|_{\text{Lip}(X, d^\alpha)}\} &\leq (p(\tau))^\alpha \|f + ig\|_{\text{Lip}(X, d^\alpha)} \\ &\leq 2(p(\tau))^\alpha \max\{\|f\|_{\text{Lip}(X, d^\alpha)}, \|g\|_{\text{Lip}(X, d^\alpha)}\}, \end{aligned}$$

for all $f, g \in A$. Assume that $\varphi : X \rightarrow X$ is a Lipschitz mapping on (X, d) satisfying $\varphi \circ \tau = \tau \circ \varphi$ and $f \in A$. It is easy to see $f \circ \varphi \in B$. Since

$$\begin{aligned} (\tau^*(f \circ \varphi))(x) &= \overline{(f \circ \varphi \circ \tau)}(x) \\ &= \overline{(f \circ \varphi)(\tau(x))} \\ &= \overline{((f \circ \varphi) \circ \tau)}(x) \\ &= \overline{(f \circ (\varphi \circ \tau))}(x) \\ &= \overline{(f \circ (\tau \circ \varphi))}(x) \\ &= \overline{((f \circ \tau) \circ \varphi)}(x) \\ &= \overline{(f \circ \tau)(\varphi(x))} \\ &= \overline{(\bar{f} \circ \tau)(\varphi(x))} \\ &= (\tau^*(\bar{f}))(\varphi(x)) \\ &= \bar{f}(\varphi(x)) \\ &= (f \circ \varphi)(x), \end{aligned}$$

for each $x \in X$, we deduce that

$$\tau^*(f \circ \varphi) = f \circ \varphi.$$

Therefore, $f \circ \varphi \in A$. Hence, (i) holds.

To prove (ii), suppose that $T : A \rightarrow A$ is a unital endomorphism of A . Let $T' : B \rightarrow B$ be the complex linear operator on B associated with T . It is easy to see that T' is a unital complex endomorphism of B . This implies that there exists a Lipschitz mapping φ on (X, d) such that $T'h = h \circ \varphi$ for all $h \in B$. Since A is a subset of B and $Tf = T'f$ for all $f \in A$, we deduce that $Tf = f \circ \varphi$ for all $f \in A$. Therefore, T is induced by φ .

Let $x \in X$. For each $f \in A$, we have

$$\begin{aligned}
 f((\varphi \circ \tau)(x)) &= (f \circ (\varphi \circ \tau))(x) \\
 &= ((f \circ \varphi) \circ \tau)(x) \\
 &= (f \circ \varphi)(\tau(x)) \\
 &= \overline{(f \circ \varphi)(x)} \\
 &= \overline{f(\varphi(x))} \\
 &= (f \circ \tau)(\varphi(x)) \\
 &= ((f \circ \tau) \circ \varphi)(x) \\
 &= (f \circ (\tau \circ \varphi))(x) \\
 &= f((\tau \circ \varphi)(x)).
 \end{aligned}$$

This implies that $(\varphi \circ \tau)(x) = (\tau \circ \varphi)(x)$ since A separates the points of X . Hence, $\varphi \circ \tau = \tau \circ \varphi$. Therefore, (ii) holds. \square

Theorem 2.5. *Let (X, d) be a compact metric space, $\tau : X \rightarrow X$ be a Lipschitz involution on (X, d) and $A = \text{Lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1]$ or $A = \text{lip}(X, d^\alpha, \tau)$ for $\alpha \in (0, 1)$. Suppose that $T : A \rightarrow A$ is a unital endomorphism of A induced by the Lipschitz mapping φ on (X, d) . Then the following statements hold.*

- (i) *If $p(\varphi_{n_0}) < 1$ for some $n_0 \in \mathbb{N}$ and $\alpha \in (0, 1)$, then $r_e(T) = \lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{\alpha}{n}}$ and T is quasicompact.*
- (ii) *If $p(\varphi_{n_0}) < 1$ for some $n_0 \in \mathbb{N}$ and $\alpha = 1$, then $r_e(T) \leq \lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{\alpha}{n}}$ and T is quasicompact.*
- (iii) *If (X, d) is connected and T is quasicompact, then $p(\varphi_{n_0}) < 1$ for some $n_0 \in \mathbb{N}$.*
- (iv) *If $\lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{1}{n}} = 0$, then T is Riesz.*
- (v) *If (X, d) is connected, $\alpha \in (0, 1)$ and T is Riesz, then $\lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{1}{n}} = 0$.*
- (vi) *T is quasicompact if and only if there exists a decomposition of X into a finite number of mutually disjoint clopen subsets, say X_1, X_2, \dots, X_m such that, for each $i \in \{1, \dots, m\}$, there exist $j \in \{1, \dots, m\}$ and $n_i \in \mathbb{N}$ with $\tau(X_i) = X_j$, $\varphi_{n_i}(X_i) \subseteq X_i$ and $p(\varphi_{n_i}|_{X_i}) < 1$.*
- (vii) *If $\alpha \in (0, 1)$ and T is quasicompact then*

$$\begin{aligned}
 \sigma_p(T) &\subseteq \{\lambda \in \mathbb{R} : |\lambda| \leq r_e(T)\} \cup \{-1, 1\}, \\
 \sigma(T) &\subseteq \{\lambda \in \mathbb{R} : |\lambda| \leq r_e(T)\} \cup \{-1, 1\}.
 \end{aligned}$$
- (viii) *If $\alpha \in (0, 1)$ and T is Riesz, then $\sigma(T) \subseteq \{0, -1, 1\}$.*

(ix) If $\alpha \in (0, 1)$, (X, d) is connected and T is Riesz, then $\sigma(T) = \{0, 1\}$.

Proof. Suppose that $B = \text{Lip}(X, d^\alpha)$ if $A = \text{Lip}(X, d^\alpha, \tau)$ and $B = \text{lip}(X, d^\alpha)$ if $A = \text{lip}(X, d^\alpha, \tau)$. By Theorem 1.1, B is the complexification of A under the injective real algebra homomorphism $J : A \rightarrow B$ defined by $J(f) = f$ ($f \in A$) and $\|\cdot\|_{\text{Lip}(X, d^\alpha)}$ is a norm on B satisfying

$$\begin{aligned} \max\{\|f\|_{\text{Lip}(X, d^\alpha)}, \|g\|_{\text{Lip}(X, d^\alpha)}\} &\leq (p(\tau))^\alpha \|f + ig\|_{\text{Lip}(X, d^\alpha)} \\ &\leq 2(p(\tau))^\alpha \max\{\|f\|_{\text{Lip}(X, d^\alpha)}, \|g\|_{\text{Lip}(X, d^\alpha)}\}, \end{aligned}$$

for all $f, g \in A$. Let $T' : B \rightarrow B$ be the complex linear operator on B associated with T . It is easy to see that T' is a unital endomorphism of B which is induced by the Lipschitz mapping φ on (X, d) . Thus, by Theorem 2.2, we have

$$(2.3) \quad \sigma_p(T) = \mathbb{R} \cap \sigma_p(T'),$$

$$(2.4) \quad \sigma(T) = \mathbb{R} \cap \sigma(T').$$

Moreover, we deduce that

$$(2.5) \quad r_e(T') = r_e(T),$$

T is quasicompact if and only if T' is quasicompact and T is Riesz if and only if T' is Riesz by Theorem 2.3.

Suppose that $p(\varphi_{n_0}) < 1$ for some $n_0 \in \mathbb{N}$ and $\alpha \in (0, 1)$. Then $r_e(T') = \lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{\alpha}{n}}$ by [7, Theorem 2.2] and T' is quasicompact by part (i) of [7, Corollary 2.4]. Therefore, $r_e(T) = \lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{\alpha}{n}}$ and T is quasicompact. Thus, (i) holds.

Let $p(\varphi_{n_0}) < 1$ for some $n_0 \in \mathbb{N}$ and $\alpha = 1$. Then

$$r_e(T') \leq \lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{\alpha}{n}},$$

by [7, Theorem 2.3] and T' is quasicompact by part (i) of [7, Corollary 2.4]. Therefore,

$$r_e(T) \leq \lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{\alpha}{n}},$$

and T is quasicompact. Thus, (ii) holds.

Suppose that (X, d) is connected and T is quasicompact. Then T' is quasicompact. Hence, $p(\varphi_{n_0}) < 1$ for some $n_0 \in \mathbb{N}$ by part (ii) of [7, Corollary 2.4]. Thus (iii) holds.

Let $\lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{1}{n}} = 0$. By part (i) of [7, Corollary 2.5], T' is Riesz. Therefore, T is Riesz and so (iv) holds.

Suppose that (X, d) is connected, $\alpha \in (0, 1)$ and T is Riesz. Then T' is Riesz. Hence, $\lim_{n \rightarrow \infty} (p(\varphi_n))^{\frac{1}{n}} = 0$ by part (ii) of [7, Corollary 2.5] and so (v) holds.

Since $\tau : X \rightarrow X$ is a homeomorphism, (vi) holds by [7, Theorem 2.10].

Suppose that $\alpha \in (0, 1)$ and T is quasicompact. Then T' is quasicompact. Hence,

$$\begin{aligned}\sigma_p(T') &\subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq r_e(T')\} \cup \{\lambda \in \mathbb{C} : \lambda^n = 1\}, \\ \sigma(T') &\subseteq \{\lambda \in \mathbb{C} : |\lambda| \leq r_e(T')\} \cup \{\lambda \in \mathbb{C} : \lambda^n = 1\},\end{aligned}$$

for some $n \in \mathbb{N}$ by [7, Theorem 2.12]. Thus, by (2.3), (2.4) and (2.5) we get

$$\begin{aligned}\sigma_p(T) &\subseteq \{\lambda \in \mathbb{R} : |\lambda| \leq r_e(T)\} \cup \{-1, 1\}, \\ \sigma(T) &\subseteq \{\lambda \in \mathbb{R} : |\lambda| \leq r_e(T)\} \cup \{-1, 1\}.\end{aligned}$$

Therefore, (vii) holds.

Suppose that $\alpha \in (0, 1)$ and T is Riesz. Then T' is Riesz and so $\sigma(T') \subseteq \{0\} \cup \{\lambda \in \mathbb{C} : \lambda^n = 1\}$ for some $n \in \mathbb{N}$ by part (i) of [7, Corollary 2.13]. Hence, $\sigma(T) \subseteq \{0, -1, 1\}$ by (2.4) and so (viii) holds.

Suppose that $\alpha \in (0, 1)$, (X, d) is connected and T is Riesz. Then T' is Riesz. Hence, $\sigma(T') = \{0, 1\}$ by considered remark of the end paragraph in [7]. Thus $\sigma(T) = \{0, 1\}$ by (2.4) and so (ix) holds. \square

The following result is a modification of [2, Theorem 2.4] and [2, Theorem 2.9] and shows that the class of real Lipschitz algebras of complex-valued functions is larger than the class of complex Lipschitz algebras regarded as real Banach algebras.

Theorem 2.6. *Let (X, d) be a compact metric space, $Y = X \times \{0, 1\}$ and ρ be the metric on Y defined by $\rho((x_1, j_1), (x_2, j_2)) = \max\{d(x_1, x_2), |j_1 - j_2|\}$. Suppose that $\tau : Y \rightarrow Y$ be the self-map on Y defined by*

$$\tau(x, 0) = (x, 1), \quad \tau(x, 1) = (x, 0), \quad (x \in X).$$

Then the following statements hold.

- (i) τ is a Lipschitz involution on the compact metric space (Y, ρ) and $p(\tau) = 1$.

- (ii) For $\alpha \in (0, 1]$, the map $\Lambda : \text{Lip}(X, d^\alpha) \rightarrow \text{Lip}(X, \rho^\alpha, \tau)$ defined by

$$\begin{aligned} (\Lambda f)(x, 0) &= f(x), & (f \in \text{Lip}(X, d^\alpha), x \in X), \\ (\Lambda f)(x, 1) &= \overline{f(x)}, & (f \in \text{Lip}(X, d^\alpha), x \in X), \end{aligned}$$

is an injective bounded real linear operator from $(\text{Lip}(X, d^\alpha), \|\cdot\|_{\text{Lip}(X, d^\alpha)})$ regarded as a real Banach algebra onto $(\text{Lip}(Y, \rho^\alpha, \tau), \|\cdot\|_{\text{Lip}(Y, \rho^\alpha)})$ satisfying

$$\|f\|_{\text{Lip}(X, d^\alpha)} \leq \|\Lambda f\|_{\text{Lip}(Y, \rho^\alpha)} \leq 2\|f\|_{\text{Lip}(X, d^\alpha)},$$

for all $f \in \text{Lip}(X, d^\alpha)$.

- (iii) For $\alpha \in (0, 1)$, $\Lambda(\text{lip}(X, d^\alpha)) \subseteq \text{lip}(Y, \rho^\alpha, \tau)$ and the map $\Gamma = \Lambda|_{\text{lip}(X, d^\alpha)} : \text{lip}(X, d^\alpha) \rightarrow \text{lip}(Y, \rho^\alpha, \tau)$ is an injective bounded real linear operator from $(\text{lip}(X, d^\alpha), \|\cdot\|_{\text{Lip}(X, d^\alpha)})$ regarded as a real Banach algebra onto $(\text{lip}(Y, \rho^\alpha, \tau), \|\cdot\|_{\text{Lip}(Y, \rho^\alpha)})$ satisfying

$$\|f\|_{\text{Lip}(X, d^\alpha)} \leq \|\Gamma f\|_{\text{Lip}(Y, \rho^\alpha)} \leq 2\|f\|_{\text{Lip}(X, d^\alpha)},$$

for all $f \in \text{lip}(X, d^\alpha)$.

We now show that the class of quasicompact (Riesz, respectively) unital endomorphisms of real Lipschitz algebras of complex-valued functions on compact metric spaces with Lipschitz involutions is larger than the class of quasicompact (Riesz, respectively) unital complex endomorphisms of complex Lipschitz algebras.

Theorem 2.7. *Let (X, d) be a compact metric space, $B = \text{Lip}(X, d^\alpha)$ for $\alpha \in (0, 1]$ or $B = \text{lip}(X, d^\alpha)$ for $\alpha \in (0, 1)$ and $T : B \rightarrow B$ be a unital complex endomorphism of B induced by the Lipschitz mapping φ on (X, d) . Let $Y = X \times \{0, 1\}$, ρ be the metric on Y defined by $\rho((x_1, j_1), (x_2, j_2)) = \max\{d(x_1, x_2), |j_1 - j_2|\}$ and τ be the Lipschitz involution on (Y, ρ) defined by*

$$\tau(x, 0) = (x, 1), \quad \tau(x, 1) = (x, 0) \quad (x \in X).$$

Suppose that $A = \text{Lip}(Y, \rho^\alpha, \tau)$ if $B = \text{Lip}(X, d^\alpha)$ and $A = \text{lip}(Y, \rho^\alpha, \tau)$ if $B = \text{lip}(X, d^\alpha)$. Let $\psi : Y \rightarrow Y$ be the self-map of Y defined by

$$\psi(x, 0) = (\varphi(x), 0), \quad \psi(x, 1) = (\varphi(x), 1) \quad (x \in X).$$

Then the following statements hold.

- (i) ψ is a Lipschitz involution on (Y, ρ) and $\psi \circ \tau = \tau \circ \psi$.
- (ii) If $S : A \rightarrow A$ is the composition endomorphism of A induced by ψ , then $r_e(S) = r_e(T)$ and so S is quasicompact (Riesz, respectively) if and only if T is quasicompact (Riesz, respectively).

Proof. Clearly, (i) holds. We prove (ii) in the case $B = \text{Lip}(X, d^\alpha)$ and $A = \text{Lip}(Y, \rho^\alpha, \tau)$ for $\alpha \in (0, 1]$. Define the map $\Lambda : B \rightarrow A$ by

$$\begin{aligned} (\Lambda f)(x, 0) &= f(x), & (f \in B, x \in X), \\ (\Lambda f)(x, 1) &= \overline{f(x)}, & (f \in B, x \in X). \end{aligned}$$

By Theorem 2.6, Λ is an injective bounded real linear operator from $(B, \|\cdot\|_{\text{Lip}(X, d^\alpha)})$, regarded as a real Banach algebra, onto $(A, \|\cdot\|_{\text{Lip}(Y, \rho^\alpha)})$. We can easily show that

$$\Lambda \circ T^n \circ \Lambda^{-1} = S^n,$$

for all $n \in \mathbb{N}$. Let $K \in \mathcal{K}_{\mathbb{C}}(B)$. Clearly, $K \in \mathcal{K}_{\mathbb{R}}(B)$ where B is regarded as a real Banach algebra. By open mapping theorem, Λ^{-1} is a bounded real linear operator from $(A, \|\cdot\|_{\text{Lip}(Y, \rho^\alpha)})$ into $(B, \|\cdot\|_{\text{Lip}(X, d^\alpha)})$. Hence, $\Lambda \circ K \circ \Lambda^{-1} \in \mathcal{K}_{\mathbb{R}}(A)$. Since

$$\begin{aligned} \|S^n - \Lambda \circ K \circ \Lambda^{-1}\| &= \|\Lambda \circ T^n \circ \Lambda^{-1} - \Lambda \circ K \circ \Lambda^{-1}\| \\ &= \|\Lambda \circ (T^n - K) \circ \Lambda^{-1}\| \\ &\leq \|\Lambda\| \|T^n - K\| \|\Lambda^{-1}\|, \end{aligned}$$

for all $n \in \mathbb{N}$, we deduce that

$$\|S^n\|_e \leq \|\Lambda\| \|T^n\|_e \|\Lambda^{-1}\|,$$

for all $n \in \mathbb{N}$. This implies that

$$(2.6) \quad r_e(S) \leq r_e(T).$$

Similarly, we can show that the converse of the inequality (2.6) holds. Therefore, $r_e(S) = r_e(T)$ and so (ii) holds.

To prove (ii) in the case $B = \text{lip}(X, d^\alpha)$ and $A = \text{lip}(Y, \rho^\alpha, \tau)$ for $\alpha \in (0, 1)$, it is sufficient to apply $\Gamma = \Lambda|_{\text{lip}(X, d^\alpha)}$ instead of Λ . \square

REFERENCES

1. D. Alimohammadi and A. Ebadian, *Hedberg's theorem in real Lipschitz algebras*, Indian J. Pure Appl. Math., 32 (2001), pp. 1470-1493.
2. D. Alimohammadi and S. Sefidgar, *Compact composition operators on real Banach spaces of complex-valued bounded Lipschitz functions*, J. Linear Topol. Algebra, 3 (2014), pp. 85-101.
3. F. Behrouzi, *Riesz and quasi-compact endomorphisms of Lipschitz algebras*, Houst. J. Math. 36 (2010), pp. 793-802.
4. F.F. Bonsall and J. Duncan, *Complete Normed Algebras*, Springer Verlag, New York, 1973.
5. J.F. Feinstein and H. Kamowitz, *Quasicompact and Riesz endomorphisms of Banach algebras*, J. Funct. Anal., 225 (2005), pp. 427-438.
6. J.F. Feinstein and H. Kamowitz, *Quasicompact endomorphisms of commutative semiprime Banach algebras*, Banach Center Publ., 91 (2010), pp. 159-167.

7. A. Golbaharan and H. Mahyar, *Essential spectral radius of quasicompact endomorphisms of Lipschitz algebras*, Rocky Mountain J. Math., 45 (2015), pp. 1149-1164.
8. A. Jiménez-Vargas, M. Lacruz, and M. Villegas-Vallecillos, *Essential norm of composition operators on Banach spaces of Hölder functions*, Abstr. Appl. Anal., Hindawi Publishing Corporation, (2011), Article ID 590853, 13 pages.
9. S.H. Kulkarni and B.V. Limaye, *Gleason parts of real function algebras*, Canad. J. Math., 33 (1981), pp. 181-200.
10. S.H. Kulkarni and B.V. Limaye, *Real Function Algebras*, Marcel Dekker, New York, 1992.
11. H. Mahyar, *Quasicompact and Riesz endomorphisms of infinitely differentiable Lipschitz algebras*, Southeast Asian Bull. Math., 35 (2011), pp. 249-259.
12. H. Mahyar and A.H. Sanatpour, *Quasicompact endomorphisms of Lipschitz algebras of analytic functions*, Publ. Math. Debr., 76/1-2 (2010), pp. 135-145.
13. H. Mahyar and A.H. Sanatpour, *Compact and quasicompact homomorphisms between differentiable Lipschitz algebras*, Bull. Belg. Math. Soc. Simon-Stevin, 17 (2010), pp. 485-497.
14. A.H. Sanatpour, *Quasicompact composition operators and power-contractive selfmaps*, Ann. Funct. Anal., 7 (2016), pp. 281-289.
15. D.R. Sherbert, *Banach algebras of Lipschitz functions*, Pacific J. Math., 13 (1963), pp. 1387-1399.
16. D.R. Sherbert, *The structure of ideals and point derivations of Banach algebras of Lipschitz functions*, Trans. Amer. Math. Soc., 111 (1964), pp. 240-272.

¹ DEPARTMENT OF MATHEMATICS, PAYAME NOOR UNIVERSITY, P. O. BOX: 19359-3697, TEHRAN, IRAN.

E-mail address: m_maighany@yahoo.com

² DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, ARAK UNIVERSITY, ARAK 38156-8-8349, IRAN.

E-mail address: d-alimohammadi@araku.ac.ir