

The Norm Estimates of Pre-Schwarzian Derivatives of Spirallike Functions and Uniformly Convex α -spirallike Functions

Zahra Orouji^{1*} and Rasul Aghalary²

ABSTRACT. For a constant $\alpha \in (-\frac{\pi}{2}, \frac{\pi}{2})$, we define a subclass of the spirallike functions, $SP_p(\alpha)$, the set of all functions $f \in \mathcal{A}$

$$\operatorname{Re} \left\{ e^{-i\alpha} \frac{zf'(z)}{f(z)} \right\} \geq \left| \frac{zf'(z)}{f(z)} - 1 \right|.$$

In the present paper, we shall give the estimate of the norm of the pre-Schwarzian derivative $T_f = f''/f'$ where $\|T_f\| = \sup_{z \in \Delta} (1 - |z|^2)|T_f(z)|$ for the functions in $SP_p(\alpha)$.

1. INTRODUCTION

Let \mathcal{A} denote the class of analytic functions f on the open unit disc $\Delta = \{z \in \mathbb{C} : |z| < 1\}$ normalized by

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n,$$

and let \mathcal{S} denote the subclass of \mathcal{A} consisting of univalent functions.

Let f and g be analytic in Δ . The function f is subordinate to g , written $f \prec g$ or $f(z) \prec g(z)$, if there exists an analytic function w such that $w(0) = 0$, $|w(z)| < 1$, and $f(z) = g(w(z))$ on Δ .

For a real number α with $0 \leq \alpha < 1$, a function $f \in \mathcal{A}$ is called starlike of order α if

$$\operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > \alpha, \quad z \in \Delta,$$

2010 *Mathematics Subject Classification.* 30C45, 30C80.

Key words and phrases. Pre-Schwarzian derivative, Spiral-like function, Uniformly convex function.

Received: 18 July 2017, Accepted: 06 October 2017.

* Corresponding author.

and f is called convex of order α if

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > \alpha, \quad z \in \Delta.$$

We denote by $\mathcal{S}^*(\alpha)$ and $\mathcal{K}(\alpha)$ the classes of starlike and convex functions of order α , respectively. It follows that $f(z) \in \mathcal{K}(\alpha)$ if and only if $zf'(z) \in \mathcal{S}^*(\alpha)$. The classes $\mathcal{S}^*(\alpha)$ and $\mathcal{K}(\alpha)$ were studied by many authors (see for example [2, 11]).

Let $T_f = f''/f'$ denote pre-Schwarzian derivative of f . Pre-Schwarzian derivative has several applications in the theory of Teichmüller spaces as well as in the theory of locally univalent functions. For a locally univalent holomorphic function f in Δ , a norm of T_f is defined by

$$\|T_f\| = \sup_{z \in \Delta} (1 - |z|^2) |T_f(z)|.$$

It is known that $\|T_f\| \leq 6$ for $f \in \mathcal{S}$ and conversely, for $f \in \mathcal{A}$, $\|T_f\| \leq 1$ implies $f \in \mathcal{S}$, and these bounds are sharp (see [1]). The norm estimates for typical subclasses of univalent functions are investigated by many authors such as [5, 6, 8]. The next result was improved by Yamashita [11].

Theorem 1.1. *Let $0 \leq \alpha < 1$ and $f \in \mathcal{A}$.*

- (i) *If $f \in \mathcal{S}^*(\alpha)$, then $\|T_f\| \leq 6 - 4\alpha$.*
- (ii) *If $f \in \mathcal{K}(\alpha)$, then $\|T_f\| \leq 4(1 - \alpha)$.*
- (iii) *If $|\operatorname{Arg}(zf'(z)/f(z))| < \alpha\pi/2$, then $\|T_f\| \leq M(\alpha) + 2\alpha$, where $M(\alpha)$ is given by*

$$M(\alpha) = \frac{4\alpha c(\alpha)}{(1 - \alpha)c^2(\alpha) + 1 + \alpha},$$

and $c(\alpha)$ is the unique solution of the following equation in the interval $(1, \infty)$:

$$(1 - \alpha)c^{\alpha+2} + (1 + \alpha)c^\alpha - c^2 - 1 = 0.$$

Remark 1.2. If $\|T_f\| < 2$ then f is bounded (see [5]).

Definition 1.3. The function f is uniformly convex (starlike) if for every circular arc γ contained in Δ with center $\xi \in \Delta$ the image arc $f(\gamma)$ is convex (starlike with respect to $f(\xi)$). The class of all uniformly convex (starlike) functions is denoted by $UCV(UST)$.

These classes were studied by A.W. Goodman [3, 4]. In [3, 4] it is shown that

$$f \in UCV \iff \operatorname{Re} \left\{ 1 + (z - \xi) \frac{f''(z)}{f'(z)} \right\} \geq 0, \quad z, \xi \in \Delta$$

and

$$f \in UST \iff \operatorname{Re} \left\{ \frac{(z - \xi)f'(z)}{f(z) - f(\xi)} \right\} \geq 0, \quad z, \xi \in \Delta.$$

Rønning [10] and Ma and Minda [7] have proved the following characterization for the functions in UCV .

Theorem 1.4. *Let $f \in \mathcal{A}$. Then $f \in UCV$ if and only if*

$$\operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \left| \frac{zf''(z)}{f'(z)} \right|, \quad z \in \Delta.$$

Corollary 1.5 ([10]). *A function $f \in \mathcal{A}$ is uniformly convex if and only if $zT_f(z) \in W$ for any $z \in \Delta$, where W is the domain*

$$\left\{ \omega = u + iv; v^2 < 2u + 1 \right\}.$$

The conformal map $g : \Delta \rightarrow \mathbb{C}$ is given by $g(0) = 0$ and

$$(1.1) \quad \begin{aligned} g(z) &= \frac{2}{\pi^2} \left(\log \frac{1 + \sqrt{z}}{1 - \sqrt{z}} \right)^2 \\ &= \frac{8z}{\pi^2} \left(1 + \frac{z}{3} + \frac{z^2}{5} + \frac{z^3}{7} + \dots \right)^2, \end{aligned}$$

which maps the unit disc Δ onto W (see [7], Example 1).

Therefore, $f \in \mathcal{A}$ is uniformly convex if and only if $zT_f(z)$ is subordinate to the function $g(z)$. Kim and Sugawa [5] give the sharp estimate of the norm of the pre-Schwarzian derivative for the functions in UCV as follow.

Theorem 1.6 ([5], Theorem 4.5). *If $f \in \mathcal{A}$ is uniformly convex, then we have*

$$(1.2) \quad \|T_f\| \leq h(t_2) = 0.94779\dots,$$

where

$$(1.3) \quad h(t) = \frac{8t^2 \cosh t}{\pi^2 \sinh^2 t}, \quad 0 < t < \infty,$$

assumes its maximum at the point $t = t_2 = 1.6061152\dots$, and equality occurs only when f is a rotation of the function $F \in \mathcal{A}$ determined by $T_F(z) = g(z)/z$, where $g(z)$ is given by (1.1).

Let Γ_ω be the image of an arc $\Gamma_z : z = z(t), (a \leq t \leq b)$ under the function $w = f(z)$. We say that the arc Γ_ω is convex α -spirallike if

$$\arg \left(\frac{z''(t)}{z'(t)} + \frac{z'(t)f''(z)}{f'(z)} \right),$$

lies between α and $\alpha + \pi$.

Definition 1.7. For a constant α with $|\alpha| < \pi/2$, the function f is uniformly convex α -spiral function if the image of every circular arc Γ_z with center at ξ lying in Δ is convex α -spirallike (see [9]).

The class of all uniformly convex α -spiral functions is denoted by $UCSP(\alpha)$. In particular, $UCSP(0) = UCV$. The next results were proved by Ravichandran and Selvaraj [9].

Lemma 1.8 ([9], Theorem 6). *A function $f(z) \in \mathcal{A}$ is in $UCSP(\alpha)$ if and only if*

$$\operatorname{Re} \left\{ e^{-i\alpha} \left(1 + \frac{zf''(z)}{f'(z)} \right) \right\} \geq \left| \frac{zf''(z)}{f'(z)} \right|, \quad z \in \Delta.$$

Lemma 1.9 ([9], Theorem 9). *Let $f(z) \in \mathcal{A}$ and $s(z)$ be defined by*

$$f'(z) = (s'(z))^{e^{i\alpha} \cos \alpha}, \quad z \in \Delta.$$

Then $f(z) \in UCSP(\alpha)$ if and only if $s(z) \in UCV$.

The main object of this paper, is investigating of the norm estimates of pre-Schwarzian derivative of the classes $UCSP(\alpha)$ and $SP_p(\alpha)$. Our results extend the result obtained by [5].

In the rest of the paper, we denote by K the value

$$(1.4) \quad 0.94774\dots$$

which is the maximum of the function h defined by (1.3) at the point $t_2 = 1.6061152\dots$

2. MAIN RESULTS

Now we can prove our first result.

Theorem 2.1. *Let $f \in \mathcal{A}$ be in $UCSP(\alpha)$. Then $\|T_f\| \leq K \cos \alpha$ where $K = 0.94774\dots$ is given by (1.4).*

Proof. Let $f \in \mathcal{A}$ be in $UCSP(\alpha)$ and $s(z)$ be defined by

$$(2.1) \quad f'(z) = (s'(z))^{e^{i\alpha} \cos \alpha}, \quad z \in \Delta.$$

By Lemma 1.9, $s(z) \in UCV$ and therefore by Theorem 1.6, $\|T_s\| \leq K$. Now, in view of (2.1) we have

$$\frac{f''(z)}{f'(z)} = e^{i\alpha} \cos \alpha \frac{s''(z)}{s'(z)}, \quad z \in \Delta,$$

and so,

$$\|T_f\| = |e^{i\alpha} \cos \alpha| \|T_s\| \leq K |\cos \alpha|.$$

□

The class of functions $F(z) = zf'(z), f(z) \in UCSP(\alpha)$ is a subclass of the spirallike functions and we denote it by $SP_p(\alpha)$. By Lemma 1.8, the function $f \in \mathcal{A}$ is in $SP_p(\alpha)$ if and only if

$$\operatorname{Re} \left\{ e^{-i\alpha} \frac{zf'(z)}{f(z)} \right\} \geq \left| \frac{zf'(z)}{f(z)} - 1 \right|, \quad z \in \Delta.$$

Geometrically it means that $zf'(z)/f(z)$ lies in the parabolic region

$$\Omega_\alpha = \left\{ \omega : \operatorname{Re}\{e^{-i\alpha}\omega\} > |\omega - 1| \right\}.$$

In the next theorem, we shall give the estimate for the norm of pre-Schwarzian derivative of the class $SP_p(\alpha)$.

Theorem 2.2 ([9], Theorem 7). *A function $f \in \mathcal{A}$ is in $SP_p(\alpha)$ if and only if*

$$\frac{zf'(z)}{f(z)} \prec e^{i\alpha}(\cos \alpha P(z) - i \sin \alpha)$$

where

$$P(z) = 1 + \frac{2}{\pi^2} \left(\log \frac{1 + \sqrt{z}}{1 - \sqrt{z}} \right)^2$$

is the function which maps Δ onto $\Omega_0 = \{u + iv, v^2 < 2u - 1, u > 0\}$.

Note that Ω_0 is the interior of a parabola in the right half-plane which is symmetric about the real axis and has vertex at $(1/2, 0)$.

Theorem 2.3. *Let $f \in \mathcal{A}$ be in $SP_p(\alpha)$. Then we have*

$$\begin{aligned} \|T_f\| &\leq \max_{y \in \mathbb{R}} \frac{8}{\pi} \sqrt{\frac{1 + y^2}{y^4 + 6y^2 - 8y \tan \alpha + 1 + 4 \tan^2 \alpha}} + K \cos \alpha \\ &\leq \frac{8}{\pi} + K \cos \alpha, \end{aligned}$$

where K is given by (1.4).

Proof. Let $f \in \mathcal{A}$ be in $SP_p(\alpha)$. By setting $p(z) = zf'(z)/f(z)$ we have

$$(2.2) \quad zT_f(z) = \frac{zf''(z)}{f'(z)} = \frac{zp'(z)}{p(z)} + p(z) - 1, \quad z \in \Delta.$$

By Theorem 2.2, we have $p(z) \prec q(z)$ where

$$q(z) = 1 + \frac{2e^{i\alpha} \cos \alpha}{\pi^2} \left(\log \frac{1 + \sqrt{z}}{1 - \sqrt{z}} \right)^2, \quad z \in \Delta.$$

Therefore there exists an analytic function $w : \Delta \rightarrow \Delta$ with $w(0) = 0$ such that $p(z) = q(w^2(z))$ and so

$$(2.3) \quad p(z) = 1 + \frac{2e^{i\alpha} \cos \alpha}{\pi^2} \left(\log \frac{1 + w(z)}{1 - w(z)} \right)^2, \quad z \in \Delta.$$

Differentiating logarithmically, we obtain

$$(2.4) \quad \frac{p'(z)}{p(z)} = \frac{\frac{8e^{i\alpha} \cos \alpha}{\pi^2} \log \left(\frac{1+w(z)}{1-w(z)} \right) \frac{w'(z)}{1-w^2(z)}}{1 + \frac{2e^{i\alpha} \cos \alpha}{\pi^2} \left(\log \frac{1+w(z)}{1-w(z)} \right)^2}.$$

Upon using Schwarz-Pick Lemma, we have

$$|w'(z)|/|1-w^2(z)| \leq 1/(1-|z|^2),$$

and so by using (2.2) to (2.4), for $z \in \Delta$ yields

$$(2.5) \quad (1-|z|^2)|T_f(z)| \leq \frac{\frac{8 \cos \alpha}{\pi^2} \left| \log \frac{1+w(z)}{1-w(z)} \right|}{\left| 1 + \frac{2e^{i\alpha} \cos \alpha}{\pi^2} \left(\log \frac{1+w(z)}{1-w(z)} \right)^2 \right|} + \frac{1-|z|^2}{|z|} \frac{2 \cos \alpha}{\pi^2} \left| \log \frac{1+w(z)}{1-w(z)} \right|^2.$$

Since by (1.1),

$$g(z) = \frac{2}{\pi^2} \left(\log \frac{1+\sqrt{z}}{1-\sqrt{z}} \right)^2,$$

has positive Taylor coefficient, we see that

$$|g(w^2(z))| \leq g(|w^2(z)|) \leq g(|z|).$$

Kim and Sugawa [5], proved that

$$(2.6) \quad \sup_{z \in \Delta} \frac{1-|z|^2}{|z|} \frac{2}{\pi^2} \left| \log \frac{1+w(z)}{1-w(z)} \right|^2 \leq \sup_{0 < x < 1} (1-x^2) \frac{g(x)}{x} = K,$$

where $K = 0.94774\dots$ is given by (1.4). Set

$$(2.7) \quad I := \sup_{z \in \Delta} \frac{\frac{8 \cos \alpha}{\pi^2} \left| \log \frac{1+w(z)}{1-w(z)} \right|}{\left| 1 + \frac{2}{\pi^2} e^{i\alpha} \cos \alpha \left(\log \frac{1+w(z)}{1-w(z)} \right)^2 \right|} = \frac{8}{\pi\sqrt{2}} \sup_{(x,y) \in \Omega} \left(\frac{\cos^2 \alpha |x+iy|}{|1 + e^{i\alpha} \cos \alpha (x+iy)|^2} \right)^{\frac{1}{2}},$$

where

$$x+iy := \frac{2}{\pi^2} \left(\log \frac{1+w(z)}{1-w(z)} \right)^2,$$

belongs to $\Omega = \{x+iy, y^2 < 2x+1\}$ and so

$$I = \frac{8}{\pi\sqrt{2}} \sup_{(x,y) \in \Omega} \left(\frac{\cos^2 \alpha \sqrt{x^2+y^2}}{1 + \cos^2 \alpha (x^2+y^2) + 2x \cos^2 \alpha - 2y \sin \alpha \cos \alpha} \right)^{\frac{1}{2}}.$$

By using the maximem principle of subharmonic functions and setting $x = \frac{y^2-1}{2}$ we obtain

$$(2.8) \quad I = \frac{8}{\pi\sqrt{2}} \sup_{y \in \mathbb{R}} \left(\frac{\frac{1}{2} \cos^2 \alpha (1 + y^2)}{1 + \left(\frac{1}{4} \cos^2 \alpha\right) y^4 + \left(\frac{3}{2} \cos^2 \alpha\right) y^2 - 2y \sin \alpha \cos \alpha - \frac{3}{4} \cos^2 \alpha} \right)^{\frac{1}{2}} \\ = \frac{8}{\pi} \sup_{y \in \mathbb{R}} \left(\frac{1 + y^2}{y^4 + 6y^2 - 8y \tan \alpha + 1 + 4 \tan^2 \alpha} \right)^{\frac{1}{2}}.$$

Therefore by relations (2.5)-(2.8) we have

$$(2.9) \quad \|T_f\| \leq \sup_{y \in \mathbb{R}} \frac{8}{\pi} \sqrt{\frac{1 + y^2}{y^4 + 6y^2 - 8y \tan \alpha + 1 + 4 \tan^2 \alpha}} + K \cos \alpha.$$

We claim that the right side of (2.9) is bounded. Let

$$(2.10) \quad h(y, \alpha) = \frac{1 + y^2}{y^4 + 6y^2 - 8y \tan \alpha + 1 + 4 \tan^2 \alpha}, \quad y \in \mathbb{R}, |\alpha| < \frac{\pi}{2}.$$

Then $\frac{\partial h}{\partial \alpha} = 0$ if and only if

$$8(y^2 + 1)(1 + \tan^2 \alpha)(-y + \tan \alpha) = 0,$$

or if and only if $y = \tan \alpha$ and also $\frac{\partial h}{\partial y} = 0$ if and only if

$$2y(y^4 + 6y^2 - 8y \tan \alpha + 1 + 4 \tan^2 \alpha) = (y^2 + 1)(4y^3 + 12y - 8 \tan \alpha).$$

Hence $\partial h / \partial \alpha = \partial h / \partial y = 0$ if and only if $y = \tan \alpha = 0$. Also it is easy to see that $h_{\alpha\alpha}(0, 0) < 0$ and $h_{\alpha\alpha}(0, 0)h_{yy}(0, 0) - h_{\alpha y}^2(0, 0)$ is positive. So the function $h(y, \alpha)$ takes its maximum value at the point $y = \alpha = 0$. But in view of (2.10), we have $h(0, 0) = 1$ and so its maximum is 1.

Now, the relation (2.9) yields

$$\|T_f\| \leq \frac{8}{\pi} + K \cos \alpha$$

and the proof is complete. \square

REFERENCES

1. J. Becker and CH. Pommerenke, *Schlichtheitskriterien und Jordangebiete*, J. Reine Angew. Math., 354 (1984), pp. 74-94.
2. P.L. Duren, *Univalent Functions*, Grundlehren Math. Wiss., 259 (1983), Springer-Verlay. New York.
3. A.W. Goodman, *On uniformly convex functions*, Ann. Pol. Math., 57 (1991), pp. 87-92.
4. A.W. Goodman, *On uniformly convex functions*, J. Math. Anal. Appl., 155 (1991), pp. 364-370.

5. Y.C. Kim and T. Sugawa, *Growth and coefficient estimates for uniformly locally univalent functions on the unit disk*, Rocky Mountain J. Math., 32 (2002), pp. 179-200.
6. Y.C. Kim and T. Sugawa, *Norm estimates of the pre-Schwarzian derivative for certain classes of univalent functions*, Proc. Edinburgh Math. Soc., 49 (2006), pp. 131-143.
7. W. Ma and D. Minda, *Uniformly convex functions*, Ann. Pol. Math., 57 (1992), pp. 165-175.
8. Y. Okuyama, *The norm estimates of pre-Schwarzian derivatives of spiral-like functions*, Complex Variables Theory Appl., 42 (2000), pp. 225-239.
9. V. Ravichandran, C. Selvaraj, and R. Rajagopal, *On uniformly convex spiral functions and uniformly spirallike functions*, Soochow J. Math., 29 (2003), pp. 393-405.
10. F. Rønning, *Uniformly convex functions and a corresponding class of starlike functions*, Proc. Amer. Math. Soc., 118 (1993), pp. 189-196.
11. S. Yamashita, *Norm estimates for function starlike or convex of order alpha*, Hokkaido Math., 28 (1999), pp. 217-230.

¹ DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN.

E-mail address: z.rouji@urmia.ac.ir

² DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN.

E-mail address: raghalary@yahoo.com