

# Integral Operators on the Besov Spaces and Subclasses of Univalent Functions

Zahra Orouji and Ali Ebadian

**Sahand Communications in  
Mathematical Analysis**

Print ISSN: 2322-5807  
Online ISSN: 2423-3900  
Volume: 17  
Number: 4  
Pages: 61-69

Sahand Commun. Math. Anal.  
DOI: 10.22130/scma.2019.109347.625

Volume 17, No. 4, November 2020

Print ISSN 2322-5807  
Online ISSN 2423-3900

Sahand Communications  
in  
Mathematical Analysis



Photo by Farhad Mansoury

Sahand Mountain, Maragheh, Iran.

SCMA, P. O. Box 55181-83111, Maragheh, Iran  
<http://scma.maragheh.ac.ir>

## Integral Operators on the Besov Spaces and Subclasses of Univalent Functions

Zahra Orouji<sup>1\*</sup> and Ali Ebadian<sup>2</sup>

---

ABSTRACT. In this note, we study the integral operators  $I_g^{\gamma,\alpha}$  and  $J_g^{\gamma,\alpha}$  of an analytic function  $g$  on convex and starlike functions of a complex order. Then, we investigate the same operators on  $H^\infty$  and Besov spaces.

---

### 1. INTRODUCTION

Let  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  be the unit disk in the plane  $\mathbb{C}$ , and  $H(\mathbb{D}) := \{g : \mathbb{D} \rightarrow \mathbb{C} \mid g \text{ is analytic}\}$ . Also, let  $A$  be the subclass of  $H(\mathbb{D})$ , which its elements are of the form

$$g(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

Suppose that  $S^*(\alpha)$  is the famous subclass of  $A$ , which is starlike of order  $\alpha$  ( $0 \leq \alpha < 1$ ). Indeed,  $g \in S^*(\alpha)$  is equivalent to  $\operatorname{Re}(zg'(z)/g(z)) > \alpha$  in  $\mathbb{D}$ . Similarly, we have  $g \in K(\alpha)$  if and only if

$$\operatorname{Re} \left( 1 + \frac{zg''(z)}{g'(z)} \right) > \alpha, \quad (z \in \mathbb{D}),$$

where  $K(\alpha)$  is the subclass of  $A$  contained in the convex functions of order  $\alpha$ .

---

2020 *Mathematics Subject Classification.* 30C45, 30C80.

*Key words and phrases.* Integral operators, Besov spaces, Convex functions of complex order, Starlike functions of complex order, Schwarzian norm.

Received: 10 June 2019, Accepted: 22 October 2019.

\* Corresponding author.

As usual, we write  $S^* = S^*(0)$  and  $K = K(0)$ . For  $0 \neq b \in \mathbb{C}$ , the subclasses of  $A$ ,  $S_b^*$  and  $K_b$ , are defined by

$$S_b^* = \left\{ g \in A : \operatorname{Re} \left\{ 1 + \frac{1}{b} \left( \frac{zg'(z)}{g(z)} - 1 \right) \right\} > 0, \quad (z \in \mathbb{D}) \right\},$$

and

$$K_b = \left\{ g \in A : \operatorname{Re} \left\{ 1 + \frac{1}{b} \left( \frac{zg''(z)}{g'(z)} \right) \right\} > 0, \quad (z \in \mathbb{D}) \right\}.$$

Then, we can see that for  $0 \leq \alpha < 1$ ,

$$S_{1-\alpha}^* = S^*(\alpha), \quad K_{1-\alpha} = K(\alpha).$$

We refer to [3, 11, 12] for some important results.

For some real number  $\beta$  and non-zero complex number  $b$ , we introduce a subclass of  $H(\mathbb{D})$ ,  $P(\beta, b)$ , as follows:

$$P(\beta, b) := \left\{ g \in H(\mathbb{D}) : \operatorname{Re} \left\{ \frac{1}{b} \left( \frac{zg'(z)}{g(z)} \right) \right\} \geq \beta \text{ and } g(0) = 1 \right\}.$$

For example,  $\frac{1}{1-z}$  and  $\frac{1}{1+z}$  belong to  $P(-\frac{1}{2}, 1)$ .

Let  $g \in H(\mathbb{D})$  be locally univalent. Let

$$S_g(z) = \left( \frac{g''(z)}{g'(z)} \right)' - \frac{1}{2} \left( \frac{g''(z)}{g'(z)} \right)^2,$$

denote the Schwarzian derivative of  $g$ , and let

$$\|S_g\| = \sup_{z \in \mathbb{D}} (1 - |z|^2)^2 |S_g(z)|,$$

which denotes its Schwarzian norm.

If  $g \in K$  and  $h(z) = 1 + \frac{zg''(z)}{g'(z)}$ , then  $\operatorname{Re} h(z) > 0$  ( $z \in \mathbb{D}$ ), so  $h$  is subordinate to  $\lambda(z) = \frac{1+z}{1-z}$ , where  $\lambda$  is the half-plan mapping. Therefore,  $h(z) = \lambda(\varphi(z))$  for some Schwarz function  $\varphi$ , and we have

$$\begin{aligned} \frac{zg''(z)}{g'(z)} &= \frac{1 + \varphi(z)}{1 - \varphi(z)} - 1 \\ &= \frac{2\varphi(z)}{1 - \varphi(z)}, \end{aligned}$$

with the notation  $\psi(z) = \frac{\varphi(z)}{z}$ , where  $\psi$  is analytic and satisfies  $|\psi(z)| \leq 1$  in  $\mathbb{D}$ . Then it can be written as follows:

$$(1.1) \quad \frac{g''(z)}{g'(z)} = \frac{2\psi(z)}{1 - z\psi(z)}.$$

Hence, the Schwarzian derivative of  $g$  can be written in the following form:

$$\begin{aligned} S_g(z) &= \left( \frac{g''(z)}{g'(z)} \right)' - \frac{1}{2} \left( \frac{g''(z)}{g'(z)} \right)^2 \\ &= \frac{2\psi'(z)}{(1 - z\psi(z))^2}. \end{aligned}$$

Then, we obtain

$$(1.2) \quad |S_g(z)| \leq \frac{2}{(1 - |z|^2)^2}.$$

By the inequality (1.2), the Schwarzian norm  $\|S_g\|$  of the convex mapping is not greater than 2. If the convex mapping  $g$  is bounded, then  $\|S_g\| < 2$  (see [10]).

Finally, let  $g \in H(\mathbb{D})$ . We consider two integral operators on  $H(\mathbb{D})$ , as follows:

$$I_g^{\gamma, \alpha}(h)(z) = \int_0^z h'(w) g^\gamma(w) w^{\alpha-1} dw, \quad (z \in \mathbb{D}),$$

and

$$J_g^{\gamma, \alpha}(h)(z) = \int_0^z h(w) (g'(w))^\gamma w^{\alpha-1} dw, \quad (z \in \mathbb{D}),$$

where  $\gamma, \alpha > 0$ .

Integral operators play an important role in various fields (see [4, 8]). If  $\gamma = \alpha = 1$ , then  $I_g^{1,1}(h) = I_g(h)$  and  $J_g^{1,1}(h) = J_g(h)$ , which are Alexander operators. These integral operators have been investigated by many authors [7, 9, 13, 14].

In this note, we study  $I_g^{\gamma, \alpha}$  and  $J_g^{\gamma, \alpha}$  operators on  $K$ ,  $K(\alpha)$  and  $S_b^*$ . Here, we obtain the necessary and sufficient conditions such that  $I_g^{\gamma, \alpha}(\mathbb{D})$  and  $J_g^{\gamma, \alpha}(\mathbb{D})$  are bounded, Furthermore, we obtain the sufficient conditions such that  $|S_{I_g^{\gamma, \alpha}}| < 2$ .

## 2. INTEGRAL OPERATORS ON $K(\alpha)$ AND $S^*(\alpha)$

Now, we verify the integral operators,  $I_g^{\gamma, \alpha}$  and  $J_g^{\gamma, \alpha}$ , on  $K(\alpha)$  and  $S^*(\alpha)$ .

- Lemma 2.1.** (i) Let  $\alpha > 0, \gamma > 0, \beta \geq 0$  and  $0 \neq b \in \mathbb{C}$ , where  $(\alpha - 1)\text{Re}b \geq 0$ . If  $g \in P(\beta, b)$ , then  $I_g^{\gamma, \alpha}$  is an operator on  $K_b$ .  
 (ii) Let  $\gamma > 0, 0 \leq \alpha < 1$  and  $\beta \in \mathbb{R}$ , where  $1 \leq 2\alpha + \beta\gamma < 2$ . If  $g \in P(\beta, 1)$ , then  $I_g^{\gamma, \alpha}$  is an operator from  $K(\alpha)$  to  $K(2\alpha + \beta\gamma - 1)$ .

*Proof.* (i) Let  $h \in K_b$ . Then, for  $z \in \mathbb{D}$ ,

$$\begin{aligned}
(2.1) \quad & \operatorname{Re} \left\{ 1 + \frac{1}{b} \frac{z(I_g^{\gamma, \alpha} h)''(z)}{(I_g^{\gamma, \alpha} h)'(z)} \right\} \\
&= \operatorname{Re} \left\{ 1 + \frac{1}{b} \frac{z(z^{\alpha-1} h''(z) g^\gamma(z) + \gamma z^{\alpha-1} h'(z) g'(z) g^{\gamma-1}(z))}{z^{\alpha-1} h'(z) g^\gamma(z)} \right\} \\
&\quad + \operatorname{Re} \left\{ \frac{1}{b} \frac{z(\alpha-1) z^{\alpha-2} h'(z) g^\gamma(z)}{z^{\alpha-1} h'(z) g^\gamma(z)} \right\} \\
&= (\alpha-1) \operatorname{Re} \left( \frac{1}{b} \right) + \operatorname{Re} \left\{ 1 + \frac{1}{b} \left( \frac{z h''(z)}{h'(z)} \right) \right\} + \gamma \operatorname{Re} \left\{ \frac{1}{b} \left( \frac{z g'(z)}{g(z)} \right) \right\}.
\end{aligned}$$

By this hypothesis and (2.1), we obtain

$$\operatorname{Re} \left\{ 1 + \frac{1}{b} \left( \frac{z(I_g^{\gamma, \alpha} h)''(z)}{(I_g^{\gamma, \alpha} h)'(z)} \right) \right\} > 0.$$

Therefore,  $I_g^{\gamma, \alpha} h \in K_b$  for all  $h \in K_b$ .

(ii) We can prove this part in a similar manner as the proof of part (i). □

**Lemma 2.2.** *Let  $\alpha > 0$ ,  $0 \neq b \in \mathbb{C}$  and  $0 < \gamma < \alpha \operatorname{Re}(\frac{1}{b})$ . If  $g \in K_b$ , then  $J_g^{\gamma, \alpha}$  is an operator from  $S_b^*$  to  $K_b$ .*

*Proof.* Let  $h \in S_b^*$ , then for  $z \in \mathbb{D}$ ,

$$\begin{aligned}
& \operatorname{Re} \left\{ 1 + \frac{1}{b} \frac{z(J_g^{\gamma, \alpha} h)''(z)}{(J_g^{\gamma, \alpha} h)'(z)} \right\} \\
&= \operatorname{Re} \left\{ 1 + \frac{1}{b} \frac{z(z^{\alpha-1} h'(z) (g'(z))^\gamma + \gamma z^{\alpha-1} h(z) g''(z) (g'(z))^{\alpha-1})}{z^{\alpha-1} h(z) (g'(z))^\gamma} \right\} \\
&\quad + \operatorname{Re} \left\{ \frac{1}{b} \frac{z(\alpha-1) z^{\alpha-2} h(z) (g'(z))^\gamma}{z^{\alpha-1} h(z) (g'(z))^\gamma} \right\} \\
&= \operatorname{Re} \left\{ 1 + \frac{1}{b} \left( \frac{z h'(z)}{h(z)} \right) + \frac{\gamma}{b} \left( \frac{z g''(z)}{g'(z)} \right) + \frac{\alpha-1}{b} \right\} \\
&= \operatorname{Re} \left\{ 1 + \frac{1}{b} \left( \frac{z h'(z)}{h(z)} - 1 \right) \right\} + \gamma \operatorname{Re} \left\{ 1 + \frac{1}{b} \left( \frac{z g''(z)}{g'(z)} \right) \right\} + \alpha \operatorname{Re} \left( \frac{1}{b} \right) - \gamma.
\end{aligned}$$

By this hypothesis, we obtain  $J_g^{\gamma, \alpha} h \in K_b$ . □

**Theorem 2.3.** *Let  $\gamma > 0$ ,  $0 \leq \alpha < 1$  and  $\beta \in \mathbb{R}$ , where  $1 \leq 2\alpha + \beta\gamma < 2$ . Also, let  $g \in P(\beta, 1)$  and  $h \in K(\alpha)$ . Then, the image  $(I_g^{\gamma, \alpha} h)(\mathbb{D})$  is bounded if and only if*

$$\limsup_{|z| \rightarrow 1} (1 - |z|) \left| \frac{z h''(z)}{h'(z)} + \frac{\gamma z g'(z)}{g(z)} + \alpha + 1 \right| < 1.$$

*Proof.* By using Part (2) of Lemma 2.1, we have  $I_g^{\gamma, \alpha} h \in K$ . By replacing  $g = I_g^{\gamma, \alpha} h$  in (1.1), then there exists  $\psi \in H(\mathbb{D})$  such that

$$\frac{(I_g^{\gamma, \alpha} h)''(z)}{(I_g^{\gamma, \alpha} h)'(z)} = \frac{2\psi(z)}{1 - z\psi(z)}.$$

Therefore

$$\begin{aligned} \psi(z) &= \frac{\frac{(I_g^{\gamma, \alpha} h)''(z)}{(I_g^{\gamma, \alpha} h)'(z)}}{2 + \frac{z(I_g^{\gamma, \alpha} h)''(z)}{(I_g^{\gamma, \alpha} h)'(z)}} \\ &= \frac{\frac{h''(z)}{h'(z)} + \frac{\gamma g'(z)}{g(z)} + \frac{\alpha-1}{z}}{1 + \alpha + \frac{zh''(z)}{h'(z)} + \frac{\gamma zg'(z)}{g(z)}}. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} (2.2) \quad \frac{1 - |z|}{|1 - z\psi(z)|} &= \frac{1 - |z|}{\frac{2}{\left| \frac{zh''(z)}{h'(z)} + \frac{\gamma zg'(z)}{g(z)} + \alpha + 1 \right|}} \\ &= \frac{1}{2} (1 - |z|) \left| 1 + \alpha + \frac{zh''(z)}{h'(z)} + \frac{\gamma zg'(z)}{g(z)} \right|. \end{aligned}$$

By Theorem 2 in [10], we can conclude that the image  $(I_g^{\gamma, \alpha} h)(\mathbb{D})$  is bounded if and only if

$$\limsup_{|z| \rightarrow 1} \frac{1 - |z|}{|1 - z\psi(z)|} < \frac{1}{2},$$

and so by (2.2), the proof is complete.  $\square$

Similarly, by using Lemma 2.2, the following theorem is achieved:

**Theorem 2.4.** *Let  $0 < \gamma < \alpha$ ,  $g \in K$  and  $h \in S^*$ . Then, the image  $(J_g^{\gamma, \alpha} h)(\mathbb{D})$  is bounded if and only if*

$$\limsup_{|z| \rightarrow 1} (1 - |z|) \left| \frac{zh'(z)}{h(z)} + \frac{\gamma zg''(z)}{g'(z)} + \alpha + 1 \right| < 1.$$

By using part 2 of Lemma 2.1, we can obtain the below result:

**Corollary 2.5.** *Let  $\gamma > 0$ ,  $0 \leq \alpha < 1$  and  $\beta \in \mathbb{R}$ , where  $1 \leq 2\alpha + \beta\gamma < 2$ . If  $g \in P(\beta, 1)$  and  $h \in K(\alpha)$ , then  $\|S_{I_g^{\gamma, \alpha} h}\| \leq 2$ .*

And also, by Theorem 2.3, the below conclusion is gained:

**Corollary 2.6.** *Let  $\gamma > 0$ ,  $0 \leq \alpha < 1$  and  $\beta \in \mathbb{R}$ , where  $1 \leq 2\alpha + \beta\gamma < 2$ . If*

$$\limsup_{|z| \rightarrow 1} (1 - |z|) \left| \frac{zh''(z)}{h'(z)} + \frac{\gamma zg'(z)}{g(z)} + \alpha + 1 \right| < 1,$$

*then  $\|S_{I_g^{\gamma, \alpha} h}\| < 2$ .*

Finally, a similar corollary to those above is also true for the operator  $J_f^{\gamma, \alpha} h$ .

### 3. INTEGRAL OPERATORS ON BESOV SPACES

We use  $dA(z)$  to denote the area measure of  $\mathbb{D}$  which is normalized, so the area of  $\mathbb{D}$  is 1. We have

$$\begin{aligned} dA(z) &= \frac{1}{\pi} dx dy \\ &= \frac{r}{\pi} dr d\theta, \end{aligned}$$

where  $z = x + iy = re^{i\theta}$  and we set

$$dA_\alpha(z) = (\alpha + 1)(1 - |z|^2)^\alpha dA(z),$$

where  $\alpha > -1$ . It is clear that if  $\alpha$  is a real number then

$$\int_{\mathbb{D}} (1 - |z|^2)^\alpha dA(z) < \infty,$$

if and only if  $\alpha > -1$ .

For  $1 < p < \infty$  and  $\delta \geq 1$ , the Besov space  $B_\delta^p$  is defined as the set of all  $g \in H(\mathbb{D})$  such that

$$\begin{aligned} \|g\|_{B_\delta^p} &:= |g(0)| + \left\{ (p-1) \int_{\mathbb{D}} (1 - |z|^2)^{p-2} |g'(z)|^p dA_\delta(z) \right\}^{\frac{1}{p}} \\ &< \infty. \end{aligned}$$

For simplicity, the space  $B_1^p$  will be denoted by  $B^p$ . Many authors have studied the properties of the Besov spaces [1, 2]. The space  $H^\infty$  consists of bounded analytic functions  $g$  in  $\mathbb{D}$  where

$$\|g\|_{H^\infty} := \lim_{r \rightarrow 1^-} (\max_{|z| \leq r} |g(z)|) < \infty.$$

In this section, we study two operators  $I_g^{\gamma, \alpha}$  and  $J_g^{\gamma, \alpha}$  on  $H^\infty$  and Besov space  $B_\delta^p$ .

**Theorem 3.1.** *Let  $1 < p < \infty$  and  $\delta \geq 1$ . If  $g \in B_\delta^p$  then  $J_g^{\gamma, \alpha}$  is bounded on  $H^\infty$  and  $\|J_g^{\gamma, \alpha}\|_{B_\delta^p} \leq \|g\|_{B_\delta^p}$  where  $\gamma \leq 1$  and  $\alpha \geq 1$ .*

*Proof.* Let  $\|h\|_{H^\infty} = 1$ . Therefore,

$$\begin{aligned} \|J_g^{\gamma, \alpha}\|_{B_\delta^p}^p &= (p-1) \int_{\mathbb{D}} (1 - |z|^2)^{p-2} |h(z)(g'(z))^\gamma z^{\alpha-1}|^p dA_\delta(z) \\ &\leq (p-1) \|h\|_{H^\infty}^p \int_{\mathbb{D}} (1 - |z|^2)^{p-2} |(g'(z))^\gamma|^p dA_\delta(z) \\ &\leq \left( \|g\|_{B_\delta^p} - |g(0)| \right)^p < \infty, \end{aligned}$$

since  $\gamma \leq 1$  and  $\alpha \geq 1$ . □

**Theorem 3.2.** *Let  $1 < p < \infty$ ,  $\delta \geq 1$  and  $g \in H^\infty$ . Then  $I_g^{\gamma, \alpha} \in B_\delta^p$ , where  $\alpha + \gamma \geq 1$ . Moreover,  $\|I_g^{\gamma, \alpha}\|_{B_\delta^p} \leq \|I_g^\gamma\|_{H^\infty}$ , where  $I(z) = z^{\alpha-1}$  ( $z \in \mathbb{D}$ ).*

*Proof.* Suppose that  $g \in H^\infty$ . There exists a number  $N > 0$  such that  $\frac{|g(z)|}{N} < |z|$  ( $z \in \mathbb{D}$ ), thus,

$$|z^{\alpha-1}g^\gamma(z)| \leq N^\gamma, \quad (z \in \mathbb{D}),$$

where  $\gamma + \alpha - 1 \geq 0$ .

Therefore,  $z^{\alpha-1}g^\gamma(z) \in H^\infty$ . Set  $I(z) = z^{\alpha-1}$  ( $z \in \mathbb{D}$ ), so there exists a number  $c > 0$  such that  $\|I_g^\gamma\|_{H^\infty} = c$ . Now, for any  $\|h\|_{B_\delta^p} = 1$ , we have

$$\begin{aligned} \|I_g^{\gamma, \alpha}h\|_{B_\delta^p}^p &= (p-1) \int_{\mathbb{D}} (1-|z|^2)^{p-2} |h'(z)z^{\alpha-1}g^\gamma(z)|^p dA_\delta(z) \\ &\leq c^p(p-1) \int_{\mathbb{D}} (1-|z|^2)^{p-2} |h'(z)|^p dA_\delta(z) \\ &\leq c^p \|h\|_{B_\delta^p}^p \\ &= c^p, \end{aligned}$$

and the proof is complete.  $\square$

Let  $\lambda > 0$  and  $g$  be a locally univalent function. Also let

$$B(\lambda) = \{g \in H(\mathbb{D}); \left\| \frac{g''}{g'} \right\| \leq 2\lambda\},$$

where

$$\left\| \frac{g''}{g'} \right\| = \sup_{z \in \mathbb{D}} (1-|z|^2) \left| \frac{g''(z)}{g'(z)} \right|,$$

is the norm of the pre-Schwarzian derivative  $\frac{g''}{g'}$  of  $g$ . Kim and Sugawa [5, 6] investigated the properties of the class  $B(\lambda)$ .

**Theorem 3.3.** *Let  $1 < p < \infty$ ,  $\delta > 1$  and  $\lambda < 1$ . Therefore,  $B(\lambda) \subseteq B_\delta^p$ .*

*Proof.* Let  $|z| = t < 1$  and  $g \in B(\lambda)$ . Then we have

$$\begin{aligned} \log \left| \frac{g'(z)}{g'(0)} \right| &\leq \left| \log \frac{g'(z)}{g'(0)} \right| \\ &= \left| \int_0^z \frac{g''(w)}{g'(w)} dw \right| \\ &\leq t \int_0^1 \left| \frac{g''(rz)}{g'(rz)} \right| dr \\ &\leq t \int_0^1 \frac{2\lambda}{1-t^2r^2} dr \end{aligned}$$



$$= 2\lambda \log \sqrt{\frac{1+t}{1-t}}.$$

This implies

$$(3.1) \quad |g'(z)| \leq |g'(0)| \left( \frac{1+t}{1-t} \right)^\lambda, \quad (|z| = t < 1),$$

therefore by using relationship (3.1), we can obtain

$$(3.2) \quad \begin{aligned} \sup_{z \in \mathbb{D}} (1 - |z|^2) |g'(z)| &\leq |g'(0)| \sup_{0 < t < 1} (1 - t^2) \left( \frac{1+t}{1-t} \right)^\lambda \\ &\leq 2^{1+\lambda} |g'(0)| \sup_{0 < t < 1} (1 - t)^{1-\lambda} \\ &= 2^{1+\lambda} |g'(0)|. \end{aligned}$$

If we set  $m = 2^{1+\lambda} |g'(0)|$  and use relationship (3.2), it is deduced that

$$\begin{aligned} \int_{\mathbb{D}} |g'(z)|^p (1 - |z|^2)^{p-2} dA_\delta(z) &= (\delta + 1) \int_{\mathbb{D}} |g'(z)|^p (1 - |z|^2)^{p+\delta-2} dA(z) \\ &\leq (\delta + 1) m^p \int_{\mathbb{D}} (1 - |z|^2)^{\delta-2} dA(z) < \infty, \end{aligned}$$

where  $\delta > 1$  and finally it is concluded that  $g \in B_\delta^p$ .  $\square$

**Theorem 3.4.** *Assume that  $\alpha + \gamma \geq 1$  and  $g \in H^\infty$ . Then the integral operator  $I_g^{\gamma, \alpha}$  is compact from  $B_\delta^p$  space to  $B_\delta^p$  space where  $1 < p < \infty$  and  $\delta \geq 1$ .*

*Proof.* Let  $g \in H^\infty$  and  $(h_n)$  be a sequence in  $B_\delta^p$  such that  $h_n \rightarrow 0$ . For  $n = 1, 2, 3, \dots$ , we have

$$\begin{aligned} \|I_g^{\gamma, \alpha} h_n\|_{B_\delta^p}^p &= (p-1) \int_{\mathbb{D}} (1 - |z|^2)^{p-2} |h_n'(z) g^\gamma(z) z^{\alpha-1}|^p dA_\delta(z) \\ &\leq \|g\|_{H^\infty}^p \cdot \|h_n\|_{B_\delta^p}^p. \end{aligned}$$

Since for  $h_n \rightarrow 0$  on  $\overline{\mathbb{D}}$ , we have  $\|h_n\|_{B_\delta^p} \rightarrow 0$  and by considering  $n \rightarrow \infty$  in the last inequality, we obtain that

$$\lim_{n \rightarrow \infty} \|I_g^{\gamma, \alpha} h_n\|_{B_\delta^p} = 0.$$

Therefore,  $I_g^{\gamma, \alpha}$  is compact.  $\square$

#### REFERENCES

1. J.J. Donaire, D. Girela, and D. Vukotić, *On the growth and range of functions in Möbius invariant spaces*, J. Anal. Math., 112(1) (2010), pp. 237-260.

2. J.J. Donaire, D. Girela, and D. Vukotic, *On univalent functions in some Mobius invariant spaces*, J. Reine. Angew. Math., 553 (2002), pp. 43-72.
3. A. Ebadian and J. Sokól, *Volterra type operator on the convex functions*, Hacet. J. Math. Stat., 47(1) (2018), pp. 57-67.
4. C. Hammond, *The norm of a composition operator with linear symbol acting on the Dirichlet space*, J. Math. Anal. Appl., 303(2) (2005), pp. 499-508.
5. Y.C. Kim and T. Sugawa, *Growth and coefficient estimates for uniformly locally univalent functions on the unit disk*, Rocky Mt. J. Math., 32 (2002), pp. 179-200.
6. Y.C. Kim and T. Sugawa, *Uniformly locally univalent functions and Hardy spaces*, J. Math. Anal. Appl., 353(1) (2009), pp. 61-67.
7. S. Li, *Volterra composition operators between weighted bergman spaces and bloch type spaces*, J. Korean Math. SOC., 45(1) (2008), pp. 229-248.
8. S. Li and S. Stević, *Integral type operators from mixed-norm spaces to  $\alpha$ -Bloch spaces*, Integr. Transf. Spec. F., 18(7) (2007), pp. 485-493.
9. S. Li and S. Stević, *Products of integral-type operators and composition operators between bloch-type spaces*, J. Math. Anal. Appl., 349(2) (2009), pp. 596-610.
10. Z. Nehari, *A property of convex conformal maps*, J. Anal. Math., 30(1) (1976), pp. 390-393.
11. Z. Orouji and R. Aghalary, *The norm estimates of pre-schwarzian derivatives of spirallike functions and uniformly convex alpha-spirallike functions*, Sahand Commun. Math. Anal., 12(1) (2018), pp. 89-96.
12. M. Taati, S. Moradi, and S. Najafzadeh, *Some properties and results for certain subclasses of starlike and convex functions*, Sahand Commun. Math. Anal., 7(1) (2017), pp. 1-15.
13. J. Xiao, *Holomorphic  $Q$  classes*, Lecture notes in mathematics, 2001.
14. K. Zhu, *Operator theory in function spaces*, MR 92c, 47031, 1990.

---

<sup>1</sup> DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN.

*Email address:* z.ouji@urmia.ac.ir

<sup>2</sup> DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN.

*Email address:* a.ebadian@urmia.ac.ir